FREEDOM DIAGNOSTICS

Certificate of Analysis

FreedomDiagnosticsTesting.com

Admin@FreedomDiagnostics.net



Client:	Beltrix
Accession #:	2609040605
Search Code:	Belt2609040605
Received:	09/04/2026
Reported:	9/7/2026
Lot:	KS752963

Sample Summary

Product:	Kisspeptin-10 10mg	**Purity:**	Vial 1: 99.72% Vial 2: 99.84% Vial 3: 99.90%
Identity:	Confirmed	**Net Content:**	Vial 1: 10.38 mg Vial 2: 10.14 mg Vial 3: 10.26 mg
Appearance:	White Lyophilized Powder		
Microbial Analysis (PCR):	Pass		
Elemental Impurities (ICP-MS): **As | Cd | Pb | Hg**	Pass		

Analytical Results

Test	Result
Identity (LC-MS)	Kisspeptin
Purity (HPLC-UV)	99.82%
Net Content	10.26 mg

Analytical Results

Test	Result
Elemental Impurities (ICP-MS):	Pass — Arsenic, Cadmium, Lead, and Mercury below reporting/specification limits

Method: Microbial detection performed using validated polymerase chain reaction (PCR)-based assay targeting common microbial contaminants.


Microbial Analysis (PCR) No Detectable Microbial DNA

Method: Endotoxin testing performed using Limulus Amebocyte Lysate assay in accordance with USP <85> under validated laboratory conditions.


Endotoxin Replicate 1:	Pass	**Assay Sensitivity: ≤0.05 EU/mL**
Endotoxin Replicate 2:	Pass	**Assay Sensitivity: ≤0.05 EU/mL**

Method: HPLC with UV detection coupled with mass spectrometry (LC-MS).

Chromatogram



Mass Confirmation



Alex Johnson

Principal Chemist

FreedomDiagnosticsTesting.com
Admin@FreedomDiagnostics.net
Proudly Owned and Operated in the USA

The peptide purity analysis reported here was conducted using LCMS/MS under standard laboratory conditions. This analysis is intended for informational purposes only and is specific to the sample (s) provided. The peptides tested are intended for research use only and are not approved for human or veterinary use, diagnostic, therapeutic, or clinical applications. Results should be interpreted by qualified professionals within the scope of the intended research. The accuracy and reliability of the test may be influenced by sample integrity, handling, and other experimental variables.